

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/20	A2	(11) International Publication Number: WO 97/39749 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US97/06897 (22) International Filing Date: 25 April 1997 (25.04.97) (30) Priority Data: 08/635,179 25 April 1996 (25.04.96) US 08/842,454 24 April 1997 (24.04.97) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: ABBRUZZESE, Bonnie, Chandler; 5673 Keating Drive, Dublin, OH 43017 (US). McCAMISH, Mark, Anthony; 1084 Woodman Drive, Worthington, OH 43085 (US). COPE, Frederick, Oliver; 6826 McGregor Street, Worthington, OH 43085 (US). DEMICHELE, Stephen, Joseph; 5525 Windwood Drive, Dublin, OH 43017 (US). (74) Agents: BRAINARD, Thomas, D.; Abbott Laboratories, Ross Products Division, Dept. 108140/DS-1, 625 Cleveland Avenue, Columbus, OH 43215-1724 (US) et al.		(81) Designated States: AU, CA, JP, MX, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD FOR THE PREVENTION AND TREATMENT OF CACHEXIA AND ANOREXIA (57) Abstract The present invention relates to methods and nutritional compositions for the prevention and treatment of cachexia and anorexia. The methods of the invention comprise administering a composition comprising effective amounts of ω -3 fatty acids such as alpha-linolenic acid, stearidonic acid, eicosapentaenoic acid, docosapentaenoic acid, docosahexaenoic acid or mixtures thereof; of branched-chain amino acids valine, leucine, isoleucine or mixtures thereof; with or without reduced levels of tryptophan and 5-hydroxytryptophan; and of antioxidant system selected from the group comprising beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD FOR THE PREVENTION AND TREATMENT OF CACHEXIA AND ANOREXIA

The present invention relates to methods and nutritional compositions for the prevention and treatment of cachexia and anorexia. In the practice of the present invention patients are enterally administered ω -3 fatty acids including, but not limited to alpha-linolenic (18:3 ω -3), stearidonic (18:4 ω -3), eicosapentaenoic (20:5 ω -3),
5 docosapentaenoic (22:5 ω -3), and docosahexaenoic (22:6 ω -3), in combination with antioxidants including, but not limited to, beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof; a source of amino-nitrogen with high levels of branched-chain amino acids such as valine, leucine, and isoleucine and with or without reduced levels of tryptophan and 5-hydroxytryptophan.

10

BACKGROUND

Cachexia is a syndrome characterized by anorexia, weight loss, premature satiety, asthenia, loss of lean body mass, and multiple organ dysfunction. The majority of patients with cancer whose disease progresses to metastatic disease develop
15 cachexia during their treatment program and the cachexia contributes to their deaths. The frequency of weight loss in cancer patients ranges from 40% for patients with breast cancer, acute myelocytic leukemia, and sarcoma to more than 80% in patients with carcinoma of the pancreas and stomach. About 60% of patients with carcinomas of the lung, colon or prostate have experienced weight loss prior to beginning chemotherapy.
20 Although the relationship between pretreatment malnutrition (weight loss) and adverse outcome is established, no consistent relationship has been demonstrated between the development of cachexia and tumor size, disease stage, and type or duration of the malignancy. Development of cachexia in the cancer patient is not caused simply by increased energy expenditure by the host or by the tumor. The malignant cachexia is
25 partially related to reduced caloric intake.

Cancer cachexia is not simply a local effect of the tumor. Alterations in protein, fat, and carbohydrate metabolism occur commonly. For example, abnormalities in carbohydrate metabolism include increased rates of total glucose turnover, increased hepatic gluconeogenesis, glucose intolerance and elevated glucose levels. Increased
30 lipolysis, increased free fatty acid and glycerol turnover, hyperlipidemia, and reduced

lipoprotein lipase activity are frequently noted. The weight loss associated with cancer cachexia is caused not only by a reduction in body fat stores but also by a reduction in total body protein mass, with extensive skeletal muscle wasting. Increased protein turnover and poorly regulated amino acid oxidation may also be important. Presence of
5 host-derived factors produced in response to the cancer have been implicated as causative agents of cachexia, e.g., tumor necrosis factor- α (TNF) or cachectin, interleukin-1 (IL-1), IL-6, gamma-interferon (IFN), and prostaglandins (PGs) (e.g., PGE₂).

Anorexia, with progressive depletion of body stores leading to the cachectic state, is observed in 50% of cancer-bearing patients. Different mechanisms proposed to
10 explain the pathogenesis of anorexia include: (i) increased production of cytokines such as TNF and IL-1, and (ii) increased serotonergic activity within the central nervous system secondary to enhanced availability to the brain of its precursor, tryptophan. Dickerson, J.W.T. et al., 1976, *J. Neurochem* 27: 1245-1247 have suggested that diets should be selected to keep the ratio of plasma tryptophan to the sum of neutral amino
15 acids constant. Cangiano, C., et al., 1994, *Anticancer Res.* 14: 1451-1456 has also disclosed that a close relationship between plasma free tryptophan concentration and anorexia in cancer patients supports the serotonergic system activity in the pathogenesis of cancer anorexia.

Cancer is characterized primarily by an increase in the number of abnormal cells
20 derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, and lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant metastatic sites. Clinical data and molecular biologic studies indicate that cancer is a multi-step process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia causing metabolic effects such as cachexia.

25 Tumor cells differ from normal cells in their metabolism of fat in that tumor cells consume short-chain and medium-chain fatty acids poorly. For example, tumor-bearing mice fed a diet rich in medium-chain triglycerides had less weight loss with a marked reduction in tumor size compared with animals fed long-chain triglycerides. Moreover, there have been problems reported with the use of high levels of medium-chain
30 triglycerides and use of structured lipids has been suggested in some total parenteral nutrition formulas. Moreover, these structured lipids do not provide the same benefits if administered enterally. U.S. Patent Nos. 4,906,664 and 5,081,105 disclose the use of certain structured lipids in the treatment of cancer. Preparations for enteral nourishment including varying ratios of ω -6 to ω -3 (2.1:1 - 3.0:1) have also been used in oncologic

patients. However, these preparations used proportionately larger amounts of ω -6 to ω -3 fatty acids. Furthermore, these preparations did not include additional amounts of branched-chain amino acids and antioxidants as set forth in the present invention. The use of the polyunsaturated fatty acid eicosapentaenoic acid is suggested for the

5 treatment of cachexia by inhibiting lipolytic activity of lipolytic agents in body fluids and the activity of the enzyme guanidino-benzoate. See Tisdale, M.J., and Beck, A., U.S. Patent No. 5,457,130, issued October 10, 1995; and Tisdale, et al. *Cancer Research* 50: 5022-5026 (August 1990). However, the product taught by Tisdale was in a solid dosage form, requiring an already ill patient to swallow 12-16 capsules per day. This method had

10 serious drawbacks, including difficulty in swallowing, belching, and bad odor.

Thus, the prevention and/or treatment of cachexia and anorexia remain a frustrating problem. Both animal and human studies suggest that nutritional support is largely ineffective in repleting lean body mass in the cancer-bearing host. Randomized trials exploring the usefulness of total parenteral nutrition (TPN) support as an adjunct to

15 cytotoxic antineoplastic therapy have demonstrated little improvement in treatment results. See for example Brennan, M.F., and Burt, M.E., 1981, *Cancer Treatment Reports* 65 (Suppl. 5): 67-68. This, along with a clear demonstration that TPN can stimulate tumor growth in animals suggests the routine use of TPN in cancer treatment is not justified. Kisner, D.L., 1981, *Cancer Treatment Reports* 65 (Suppl. 5): 1-2.

20 Long chain fatty acid bio-pathways and physiological actions are discussed in U.S. Patent 5,223,285 to DeMichele, et al., the entirety of which is incorporated herein by reference.

Also of interest is U.S. Patent 5,444,054 to Garleb, et al. and a related U.S. Patent _____ (allowed application Serial No. 08/221,349). These documents

25 describe compositions and methods useful in the treatment of ulcerative colitis. Such compositions include a protein source that can be intact or hydrolyzed proteins of high biological value (col. 21); an indigestible oligosaccharide such as fructooligosaccharide; and a lipid blend containing a relatively high proportion of eicosapentanoic acid, which contributes to a relatively high ω -3 to ω -6 fatty acid ratio.

30

SUMMARY OF THE INVENTION

The methods of the invention generally comprise inhibiting metabolic and cytokine associated features of cachexia in an individual by administering a nutritional composition comprising an effective amount of ω -3 fatty acids including, but not limited to

- 5 alpha-linolenic acid, stearidonic acid, eicosapentaenoic acid, docosapentaenoic acid and docosahexaenoic acid, alone or in combination with each other. The invention also relates to administering a nutritional composition comprising effective amounts of branched-chain amino acids, valine, leucine, isoleucine, or mixtures thereof, and with or without a reduced amount of tryptophan and hydroxytryptophan. The invention further
- 10 provides a method of reducing oxidative damage and anti-cancer drug-induced immunosuppression in a cancer patient by administering a nutritional composition comprising effective amounts of antioxidants including, but not limited to beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof.

- In one aspect, the invention provides a method of preventing the onset of cachexia
- 15 and/or anorexia, or treating existing cachexia and/or anorexia in a human comprising enterally administering to the human at least:

- (a) an oil blend containing ω -6 fatty acids and at least 450 mg of ω -3 fatty acids, the weight ratio of ω -6 fatty acids to ω -3 fatty acids being from about 0.1 to about 3.0; and
- 20 (b) a source of amino-nitrogen wherein 15% to 50% by weight of the amino acids of said source of amino-nitrogen are branched-chain amino acids; and
- (c) an antioxidant component comprised of at least one nutrient selected from the group comprising beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof.

- These components may be administered in a single composition or in separate
- 25 vehicles. Preferably, about 15% to about 25% of the amino-nitrogen is provided by branched-chain amino acids; most preferably, about 20%. It is also preferred that the source of amino nitrogen provides tryptophan in an amount of less than about 5.0% by weight of the total amount of the amino acids of said source of amino-nitrogen; more preferably at a level of less than 3% by weight.

- 30 In another aspect, the invention provides a method of preventing the onset of anorexia or of treating existing anorexia in a human comprising administering to the human a nutritional composition comprising amino-nitrogen wherein about 5 to 25 grams of branched-chain amino acids selected from valine, leucine, isoleucine, or mixtures thereof are present in an amount from about 15% to about 50% by weight, preferably
- 35 about 15 - 25%, of the total amount of amino-nitrogen present in said nutritional

composition, and wherein tryptophan in an amount not greater than about 5.0 % by weight of the total amount of amino acids is present in said composition and wherein ω -6 and ω -3 fatty acids are present at a weight ratio of from about 0.1 to about 3.0 and at least one antioxidant is present in the nutritional composition.

5 There is further disclosed a method for preventing immunosuppression in a human comprising administering to the human a liquid nutritional composition comprising:

- (a) an oil blend containing ω -6 and ω -3 fatty acids, the weight ratio ω -6 fatty acids to ω -3 fatty acids being about from 0.1 to about 3.0; and
 - (b) an antioxidant component comprising about 2,500 to about 6,500
- 10 micrograms per liter beta-carotene, about 250 to about 1,000 milligrams per liter vitamin C, about 100 to about 500 I.U. per liter vitamin E, and about 75 to about 125 mcg per liter selenium.

There is also disclosed a method of enhancing the transport and efficacy of anticancer drugs in a human having a cancerous condition comprising administering to

15 the human a nutritional composition comprising an oil blend containing ω -6 and ω -3 fatty acids, the weight ratio of total ω -6 fatty acids to ω -3 fatty acids being from about 0.1 to about 3.0.

In another aspect, the invention provides a liquid nutritional composition comprising per liter :

- 20 (a) at least 0.45 gm (450 mg) of ω -3 fatty acids and wherein the weight ratio of ω -6 fatty acids to ω -3 fatty acids is from about 0.1 to about 3.0;
- (b) at least 50 grams of a source of amino-nitrogen wherein 15 to 50% by weight of the amino-nitrogen is branched-chain amino acids and wherein tryptophan is present in an amount less than about 5.0% by weight of the total amino-nitrogen; and
- 25 (c) at least 1 gram of an antioxidant system comprising beta-carotene, vitamin C, vitamin E and selenium.

Generally, such compositions provide much higher levels of the ω -3 fatty acids: preferably from about 1.0 gm to about 100 gm per liter; more preferably, from about 5.0 gm to about 10 gm per liter. Similarly, it is preferred that about 15 - 25% (typically about

30 20%) by weight of the source of amino-nitrogen is branched-chain amino acids.

The various methods according to the present invention may be accomplished by feeding a single composition that contains all the components of the invention (ω -6 to ω -3 oil, branched-chain amino acids and antioxidant system) or each component may be fed individually. Further, these methods may be accomplished through the consumption of

35 pills or capsules that contain the elements of the claimed invention. In one embodiment of

the invention, a nutritional liquid formulation containing all the elements of the invention is contemplated except for the branched-chain amino acids which may be consumed in the form of a pill or tablet.

In yet another co-embodiment of the invention, a liquid nutritional product contains
5 all the elements of the composition, wherein the branched-chain amino acids are dispersed within the liquid in the form of microcapsules. This administration of the branched-chain amino acids in the form of capsules, tablets, pills and/or microcapsules is advantageous since the organoleptic or taste properties of the amino acids are very objectionable.

10 In contrast to the prior art, the nutritional composition of the present invention is not restricted to correcting metabolism of just one nutrient class at a time, such as lipids or amino acids. Instead, a preferred nutritional multinutrient composition comprises a balanced formulation containing ω -3 fatty acids, antioxidants, branched-chain amino acids, and with or without a reduced level of tryptophan and 5-hydroxytryptophan. Such a
15 composition can demonstrate strong inhibition of cachexia and anorexia associated with a variety of different cancers (disease states).

In yet another embodiment, the methods further optionally comprise administering the nutritional composition in combination with cancer chemotherapeutic agents, including but not limited to, 5-fluorouracil, mitomycin-C, adriamycin, chloroethyl nitrosoureas and
20 methotrexate, to improve the transport of the drug into the target cancer cells and ultimately the efficacy of the anticancer agent.

The Examples presented below exemplify the use according to the methods of the invention of ω -3 fatty acids, antioxidants, branched-chain amino acids with or without a reduced level of tryptophan in nutritional therapy of cachexia and anorexia in human
25 patients suffering from different cancers, including, but not limited to, liver, breast, lung, prostate, gastro-intestinal and pancreatic cancer.

DETAILED DESCRIPTION OF THE INVENTION

"Cachexia" refers to a state of general ill health and malnutrition. It is often associated with and induced by malignant cancer, and is characterized by loss of appetite, loss of body mass, especially lean body mass, and muscle wasting.

5 "Anorexia" refers simply to a loss of appetite, whether brought on by medical or psychological factors. Anorexia is often closely associated with, and generally contributes to, the cachexia seen in patients with advanced cancers.

"Fatty acids" refer to a family of carboxylic acids having a hydrocarbon chain, generally from about 12 to 24 carbons long. When unsaturated (having a double bond) at
10 least one point in the hydrocarbon chain, such fatty acids are designated by the position of the first double bond. ω -3 fatty acids have a first double bond at the third carbon from the methyl end of the chain; and include, but are not limited to, α -linolenic acid, stearidonic acid, eicosapentaenoic acid ("EPA"), docosapentaenoic acid and docosahexaenoic acid ("DHA") and the like. ω -6 fatty acids have a first double bond at
15 the sixth carbon from the methyl end of the chain; and include, but are not limited to, linoleic acid, γ -linolenic acid, arachidonic acid ("AA"), and the like. The ratio of ω -6 fatty acids to ω -3 fatty acids is simply the ratio of the total amounts (usually expressed as weight) of each type.

Branched-chain amino acids are amino acids that have a fork or branch in the side
20 chain. These include primarily those having a carbon-carbon branch, i.e. valine, leucine and isoleucine; but may also include other types of branches.

"Nutritional matrix" as used herein refers to a delivery vehicle that contains fats, amino nitrogen and carbohydrates and provides some or all of the nutritional support for a patient in the recommended daily amounts. Frequently a nutritional matrix will contain
25 vitamins, minerals, trace minerals and the like to provide balanced nutrition.

"Cytokines" as used herein refer to the causative agents of cachexia in the cancer patient, produced by the individual in response to the presence of cancer, and include, but are not limited to, tumor necrosis factor (TNF) or cachectin, interleukin-1 (IL-1), IL-6 and gamma-interferon (IFN). TNF is produced by the macrophages in response to nonspecific
30 stimuli including cancer, infection, trauma and stress. The mechanism of action in cancer cachexia involves an immune response to the tumor with the production of cytokines, which not only mediate tumor lysis but also the metabolic changes seen in cancer cachexia through specific TNF receptors and/or via the induction of other cytokine receptors.

While not intending the invention to be limited to any particular theory of operation, applicants describe below a probable mechanism. A mode of action of cytokines is mediated via interactions with receptors on the plasma membrane. This is typically defined as a "signal transduction event." In general, a cytokine receptor consists of an
5 extracellular domain, a transmembrane region spanning the phospholipid bilayer of the plasma membrane, and an intracellular domain having either enzymatic activity or binding other molecules, so that a signal is delivered inside the cell in response to the cytokine ligand interaction. The signal transduction mechanisms involve second messengers, including phospholipases, adenylate cyclase and cyclic AMP, inositol phosphates,
10 diacylglycerols and protein kinase C. More particularly, phospholipase A2 generates arachidonic acid, a precursor of dienoic prostaglandins, thromboxanes, prostacyclin and leukotrienes of the 4 series.

Cytokines such as TNF and IL-1 stimulate production of arachidonic acid metabolites which are important to their inflammatory and tissue damaging actions and
15 are responsible for immunosuppression in general, and in exacerbating some paraneoplastic conditions including metabolic changes seen in cancer cachexia.

The invention is based, in part, on a method of inhibiting signal transduction and cytokine activity using nutritional compositions comprising high levels of ω -3 fatty acids, in particular, of the long chain (e.g. 20 or more carbons) ω -3 fatty acids, eicosapentaenoic
20 (EPA) and docosahexaenoic (DHA).

Since administration of EPA and DHA results in a reduction of arachidonic acid in membrane phospholipids, such an effect not only diminishes the supply of arachidonic acid as a precursor for the dienoic eicosanoids but also inhibits their production through competitive inhibition by EPA. The cyclooxygenase and lipoxygenase metabolites of EPA
25 have attenuated activity. Furthermore, ω -3 fatty acids, alpha-linolenic and stearidonic can be converted through elongation/desaturation to EPA; and similarly, DHA can be retroconverted to EPA. Thus, the methods of invention comprise methods of inhibiting cytokine activity (e.g., TNF, IL-1) and cancer cachexia by interfering with signal transduction at the receptor level and inhibiting arachidonic acid metabolism.

30 Incorporation of ω -3 fatty acids in membrane phospholipids not only alters the activity of membrane-associated enzymes (e.g., phospholipase A2) but also alters the balance between constituent saturated and unsaturated fatty acids and regulation of membrane fluidity, facilitates the transport of anticancer drugs into the cancer cells and thus enhances the efficacy of the drugs. Alberts, A.W., et al., 1978, *Biochim. Biophys.*
35 *Acta* 509:239-250. In addition, the inhibition of arachidonic acid metabolism results in

prevention and/or reversal of immunosuppression by reducing the production of prostaglandins and leukotrienes (PGE2 and LTB4), which are immunosuppressive.

The invention also provides a method of reducing the concentration of brain tryptophan and serotonin to prevent or inhibit premature satiety and cancer cachexia and/or anorexia in a cancer patient in whom the prevention and treatment of cancer anorexia is desired by administering effective amounts of branched-chain amino acids, valine, leucine, isoleucine, or mixtures thereof, and with or without a reduced amount of tryptophan.

The methods and compositions of the invention provide a method of manipulating the concentration of brain tryptophan by: (i) increasing the branched-chain amino acids, which provide competition for tryptophan for penetration across the blood-brain barrier; and (ii) reduced levels of tryptophan and 5-hydroxytryptophan in relation to branched-chain amino acids in the nutritional composition of the invention. Such an intervention can increase appetite and thus prevent and/or treat cancer anorexia.

The methods and compositions of the invention also provide a method of reducing the risk or progression of certain symptoms of cancer, such as cancer cachexia and anorexia by administering antioxidant nutrients including, but not limited to, beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof. Epidemiological evidence indicates that a combination of beta-carotene, vitamin E and selenium can effect a reduction in cancer risk in some populations. Blot, N.J. et al., 1993, *J. Natl. Cancer Inst.* 85: 1483-1492. Furthermore, vitamin E is added to satisfy any additional requirements as a result of a higher intake of ω -3 polyunsaturated fatty acids. By the administration of the antioxidant nutrients of the invention to cancer patients having cachexia and whose immune system has been depressed on account of chemotherapy and/or oxidative burden, improvements in the nutritional status, as well as prevention and treatment of immunosuppression and cachexia can be achieved.

Nutritional support in the cancer patient can be categorized as (i) supportive, in which nutrition support is instituted to prevent nutrition deterioration in the adequately nourished patient or to rehabilitate the depleted patient before definitive therapy; (ii) adjunctive, in which nutrition support plays an integral role in the therapeutic plan; and (iii) definitive, in which aggressive nutrition support is required for the patient's existence. The routes for providing nutrition support include an oral diet, tube feeding and peripheral or total parenteral nutrition. The preferred embodiment for nutritional methods and compositions of the invention is by the oral route.

An alternate to oral feeding is tube feeding by means of nasogastric, nasoduodenal, esophagostomy, gastrostomy, or jejunostomy tubes.

A typical nutritional composition useful in the method of this invention will have a caloric distribution as follows: about 12 to 24% (target 21%) from a source of amino-nitrogen, about 40 to 65% (target 61%) from carbohydrate and about 10 to 35% (target 18%) from fat. More particularly, the oil blend may comprise approximately 30% of ω -3 fatty acids, preferably largely consisting of eicosapentaenoic acid and docosahexaenoic acid. Dietary oils used in the preparation of the nutritional composition generally contain ω -3 fatty acids in the triglyceride form and include, but are not limited to canola, medium chain triglycerides, fish, soybean, soy lecithin, corn, safflower, sunflower, high-oleic sunflower, high-oleic safflower, olive, borage, black currant, evening primrose and flaxseed oil. Table 1 sets forth both preferred amounts and ranges for an oil blend useful in the invention. Specifically, the weight ratio of ω -6 fatty acids to ω -3 fatty acids in the lipid blend according to the invention is about 0.1 to 3.0. The daily delivery of ω -3 fatty acids should be at least 450 mg and may vary depending on body weight, sex, age and medical condition of the individual. As mentioned, higher levels are desired for adult human consumption: for example, from about 0.5 to 50 gm daily, more preferably from about 2.5 to 5 gm daily.

TABLE 1: OIL BLEND
(% total weight of lipid blend)

OIL	PREFERRED	RANGE
Canola	9.3%	5.0 - 40.0%
MCT	16.2%	10.0 - 50.0%
Fish	65.0%	25.0 - 80.0%
Soybean	5.5%	3.0 - 30.0%
Soy lecithin	4.0%	2.0 - 6.0%

Table 2 presents the fatty acid profile of an exemplary oil blend useful in the present invention. The weight ratio of the total ω -6 fatty acids to the total ω -3 fatty acids in this embodiment is 0.26 to 1 which is within the claimed range for this invention.

TABLE 2: FATTY ACID PROFILE

(% of total fatty acids by weight)

OIL	%
5 Caproic (6:0)	0.53
Capyrlic (8:0)	10.35
Capric (10:0)	7.16
Lauric (12:0)	0.29
Myristic (14:0)	3.53
Palmitic (16:0)	7.41
10 Palmitoleic (16:1 ω 7)	5.73
Stearic (18:0)	1.39
Oleic (18:1 ω 9)	15.23
Linoleic (18:2 ω 6)	7.21
Gamma-linoleic (18:3 ω 6)	0.21
15 Alpha-linoleic (18:3 ω 3)	2.21
Stearidonic (18:4 ω 3)	2.40
Arachidic (20:0)	0.13
Eicosenoic (20:1 ω 9)	0.74
Arachidonic (20:4 ω 6)	0.87
20 Eicosapentaenoic (20:5 ω 3)	17.14
Erucic (22:1 ω 9)	0.17
Docosapentaenoic (22:5 ω 3)	2.08
Docosahexaenoic (22:6 ω 3)	7.73
Nervonic (24:1 ω 9)	0.14
25 Others	7.35
Total	100.00

TABLE 3: LIPID BLEND CHARACTERISTICS

30	% ω -3 fatty acids	30.51
	% ω -6 fatty acids	9.67
	% ω -9 fatty acids	15.28
35	% saturated fatty acids	27.07
	% monounsaturated fatty acids	19.33
	% polyunsaturated fatty acids	40.17
	ω -6/ ω -3 ratio	0.32
	18:2 ω 6/18:3 ω 3 ratio	4.26
40	18:3 ω 3, % total energy	0.33
	18:2 ω 6, % total energy	1.41
	18:1 ω 9, % total energy	2.42
	PUFAs, % total calories	7.23
45	saturated fatty acids, % total calories	4.87
	EPA (20:5 ω 3) per 8 oz container, g	1.09
	DHA (22:6 ω 3) per 8 oz container, g	0.46

Table 3 (above) sets forth selected characteristics of an oil blend useful in the method of this invention. However, it will be realized that the characteristics may vary among other formulas useful for this invention, depending on the specific oils added and the ratios in which they are used.

An amino acid profile for a nutritional composition useful in the invention is presented in Table 4.

TABLE 4: AMINO ACID PROFILE

	<u>Amino Acid</u>	<u>g/100g Protein</u>
10	Aspartic Acid	7.08
	Threonine	4.34
	Serine	5.68
	Glutamic Acid	20.58
	Proline	10.55
15	Glycine	1.81
	Alanine	3.04
	Valine	5.90
	Methionine	2.78
	Isoleucine	4.77
20	Leucine	9.08
	Tyrosine	4.79
	Phenylalanine	4.96
	Histidine	2.67
	Lysine	7.27
25	Arginine	3.15
	Tryptophan	0.99
	Cystine	0.56
	<u>Total BCAA</u>	<u>19.75</u>

30

The total amount of branched-chain amino acids ("BCAA") useful in the present invention is about 15-50g/100g protein (i.e. percent), preferably about 15-25g/100g. Thus, an 8 oz container of the nutritional composition would contain up to about 8g BCAAs per 16 grams of total protein. The daily delivery of BCAAs is about 5-26g. In order to deliver such a high amount of the BCAAs, and because the BCAAs impart an unpleasant taste, the nutritional composition may be accompanied by 1-3 gelatin capsules containing BCAAs to provide the additional amount required above the inherent amount present in the liquid product. The preferred BCAAs are, but are not limited to, leucine, isoleucine and valine, and are predominantly bitter in taste. Therefore, administering the additional BCAAs in encapsulated form avoids taste problems which are encountered with the use of quantities greater than 20g/100g protein of BCAAs in the liquid product. The microencapsulated BCAAs may also be mixed with taste masking compounds including,

but not limited to, polyphosphates, cyclodextrin (a cyclic glucose oligomer) and Thaumatin (a proteinaceous intense sweetener).

A representative antioxidant profile useful in the method of the invention is presented in Table 5 with range values and a preferred embodiment.

TABLE 5: ANTIOXIDANT PROFILE

	Antioxidant	Preferred	Range
10	Beta-carotene	5,000 µg/L	2,500 - 6,500 µg/L
	Vitamin E	300 IU/L	100 - 500 IU/L
	Vitamin C	650 mg/L	250 - 1,000 mg/L
	Selenium	90 µg/L	78.8 - 125 µg/L

15 The overall nutrient profile of this example is set forth in Table 6. In a specific embodiment of this invention, the nutritional product provides at least 100% of the U.S. RDA for vitamins and minerals in 1184 mL (five 8 fluid ounce servings), which would provide 1184 kcal per day.

20 If used as a sole source of nutrition, and assuming a 2000 kcal diet, between 8 and 9 servings (237 mL ; 8 fluid ounces) of this illustrative formulation would be required. However, as seen from example IV below, there is benefit derived from supplementation with as few as two servings per day. Thus a minimum daily amount of long chain ω -3 fatty acids is preferably about 3 grams, calculated as (1.06 g EPA + 0.46 g DHA) times 2
 25 8 oz .servings. Of course, if more servings are consumed to provide additional calories, more ω -3 fatty acids will be administered, up to a practical maximum of about 14 grams per day (about 9 servings at same fatty acid levels). Levels of the fatty acids, antioxidants and/or source of amino nitrogen on a per liter basis are not crucial, except to the extent
 30 that a reasonable volume of fluid should supply the recommended daily amounts consistent with the invention. Determination of a reasonable volume is easily within the ambit of those skilled in the art, especially in view of the specific guidance found in the examples.

TABLE 6: NUTRIENT PROFILE

Nutrient	Qty/Liter
Protein, g	67.40
Fat, g	27.20
Carbohydrate, g	207.00
Total Dietary Fiber, g	10.70
Indigestible Oligosaccharide (FOS), g	12.40
Gum Arabic, g	9.10
Soy Polysaccharide, g	1.60
Beta-carotene, µg	5000
Vitamin A, IU	5500
Vitamin D, IU	800.00
*Vitamin E, IU	300.00
Vitamin K, µg	135.00
Vitamin C, mg	650.00
Folic Acid µg	1900
Thiamine, mg	6.50
Riboflavin, mg	5.00
Vitamin B ₆ , mg	5.00
Vitamin B ₁₂ , µg	18.00
Niacin, mg	40.00
Choline, mg	525.00
Biotin, µg	750.00
Pantothenic Acid, mg	24.00
Sodium, mg	1500
Potassium, mg	2000
Chloride, mg	1519
Calcium, mg	1800
Phosphorous, mg	1250
Magnesium, mg	450.00
Iodine, µg	175.00
Copper, mg	2.61
Zinc, mg	29.20
Iron, mg	22.20
Selenium, µg	90.00
Chromium, µg	125.00
Molybdenum, µg	206.00
Carnitine, mg	150.00
Taurine, mg	275.00
Kcal/mL	1

* d-alpha-tocopherol (all natural form) or dl-alpha-tocopherol acetate, or a combination of the two.

5

The following specific examples are set forth to illustrate various preferred embodiments of the invention but the scope of the invention is defined by the appended claims.

10

EXAMPLE I

The specific list of materials for manufacturing the nutritional cancer product of this Example I is presented in Table 7. Of course, various changes in specific ingredients and quantities may be made without departing from the scope of the invention.

TABLE 7: LIST OF MATERIALS

<u>INGREDIENT</u>	<u>AMOUNT</u>
WATER	31,605.21 kg
GUM ARABIC	437.84 kg
ULTRATRACE/TRACE MINERAL PREMIX	14.50 kg
ZINC SULFATE	2969.89 gm
FERROUS SULFATE	2856.50 gm
MANGANESE SULFATE	784.60 gm
CUPRIC SULFATE	423.11 gm
SODIUM MOLYBDATE	21.39 gm
CHROMIUM CHLORIDE	20.80 gm
SODIUM SELENITE	8.11 gm
CITRIC ACID	894.94 gm
SUCROSE (Carrier)	6520.67 gm
POTASSIUM CITRATE	50.00 kg
SODIUM CITRATE	95.00 kg
POTASSIUM IODIDE	9.00 gm
POTASSIUM CHLORIDE	91.00 kg
CORN SYRUP SOLIDS	5630.96 kg
MALTODEXTRIN	1407.52 kg
MAGNESIUM PHOSPHATE DIBASIC	131.00 kg
CALCIUM PHOSPHATE TRIBASIC (PREFERABLY MICRONIZED)	47.50 kg
CALCIUM CARBONATE	122.50 kg
SUGAR (SUCROSE)	852.77 kg
FRUCTOOLIGOSACCHARIDE	509.96 kg
MEDIUM CHAIN TRIGLYCERIDES (FRACTIONATED COCONUT OIL)	172.69 kg
CANOLA OIL	99.13 kg
SOY OIL	58.63 kg
57% VITAMIN A PALMITATE	250.00 gm
2.5% VITAMIN D	35.00 gm
D-ALPHA-TOCOPHEROL ACETATE (R,R,R)	10.65 kg
PHYLLOQUINONE	6.50 gm
30% BETA-CAROTENE	824.00 gm
SOY LECITHIN	42.64 kg
SODIUM CASEINATE	1427.04 kg
PARTIALLY HYDROLYZED SODIUM CASEINATE	1427.04 kg
SOY POLYSACCHARIDE	85.28 kg
75% WHEY PROTEIN CONCENTRATE	184.46 kg
REFINED DEODORIZED SARDINE OIL	692.87 kg
ASCORBIC ACID	37.08 kg
45% POTASSIUM HYDROXIDE	25.96 kg
TAURINE	12.00 kg
WATER SOLUBLE VITAMIN PREMIX	4.50 kg
NIACINAMIDE	1688.60 gm
CALCIUM PANTOTHENATE	1092.24 gm
THIAMINE CHLORIDE HYDROCHLORIDE	278.78 gm

<u>INGREDIENT</u>	<u>AMOUNT</u>
PYRIDOXINE HYDROCHLORIDE	268.34 gm
RIBOFLAVIN	217.87 gm
FOLIC ACID	37.82 gm
BIOTIN	32.87 gm
CYANOCOBALAMIN	0.75 gm
DEXTROSE (Carrier)	882.74 gm
FOLIC ACID	43.50 gm
CHOLINE CHLORIDE	25.00 kg
L-CARNITINE	7.00 kg
ARTIFICIAL STRAWBERRY FLAVOR	31.75 kg
ARTIFICIAL CREAM FLAVOR	18.14 kg
FD & C Red Dye No. 3	1,220.16 gm

The liquid nutritional product of the present invention was manufactured by preparing three slurries which are blended together, combined with refined deodorized sardine oil, heat treated, standardized, packaged and sterilized. The process for manufacturing 45,360 kg (100,000 pounds) of the liquid nutritional product, using the List of Materials from Table 7, is described in detail below.

A carbohydrate/mineral slurry is prepared by first heating about 6,260 kg of water to a temperature in the range of about 71 °C to 77 °C with agitation. The gum arabic is then added to the water using a mixing apparatus. Next the ultratrace/trace mineral premix is added to the water and dissolved by agitating the resultant solution for at least one minute. The following minerals are then added, in the order listed, with high agitation: potassium citrate, sodium citrate, potassium iodide and potassium chloride. The corn syrup solids (Grain Processing Corporation, Muscatine, Iowa, U.S.A. under the trade designation "Maltrin M-200") and maltodextrin (Grain Processing Corporation, trade designation "Maltrin M-100") are then added to the slurry and the temperature of the slurry is maintained at about 71 °C with high agitation for at least about 20 minutes.

Add magnesium phosphate dibasic, calcium phosphate tribasic, and calcium carbonate to the slurry. Sugar (sucrose), and fructooligosaccharide (Golden Technologies Company, Golden, Colorado, U.S.A. under the trade designation "Nutriflora-P- Fructo-oligosaccharide Powder (96%)") are added to the slurry. The completed carbohydrate/mineral slurry is held with high agitation at a temperature in the range of about 60 - 66 °C for not longer than 12 hours until it is blended with the other slurries.

An oil slurry is prepared by combining and heating the medium chain triglycerides (fractionated coconut oil), canola oil and soy oil to a temperature in the range of about 32 - 43 °C with agitation. The 57% vitamin A palmitate, 2.5% vitamin D₃, D-alpha-tocopherol acetate (R,R,R form; Distillation Products Industries, a division of Eastman Kodak

Chemical Company, Rochester, New York U.S.A. under the trade designation "Eastman Vitamin E 6-81 D-Alpha Tocopherol Acetate Concentrate"), phylloquinone and 30% beta-carotene are added to the slurry with agitation. The soy lecithin is then added to the slurry with agitation. The completed oil slurry is held under moderate agitation at a temperature in the range of about 32 - 43 °C for not longer than 12 hours until it is blended with the other slurries.

A protein-and-fiber-in-water slurry is prepared by first heating about 19,678 kg of water to a temperature in the range of about 60 - 63 °C with agitation. Sodium caseinate, partially hydrolyzed sodium caseinate (distributed by New Zealand Milk Products, Santa Rosa, California, U.S.A. under the trade name Alanate 167) and soy polysaccharide are blended into the slurry using a mixing apparatus. The temperature of the slurry is lowered to about 57 - 60 °C and then the 75% whey protein concentrate is added to the slurry using a mixing apparatus. The completed protein-and-fiber-in-water slurry is held under agitation at a temperature in the range of about 54 - 60 °C for not longer than 2 hours before being blended with the other slurries.

The oil slurry and the protein-and-fiber-in-water slurry are blended together with agitation and the resultant blended slurry is maintained at a temperature in the range of about 54 - 66 °C. After waiting for at least one minute the carbohydrate/mineral slurry is added to the blended slurry from the preceding step with agitation and the resultant blended slurry is maintained at a temperature in the range of about 54 - 66 °C. The vessel which contained the carbohydrate/mineral slurry should be rinsed with about 220 kg of water and the rinse water should be added to the blended slurry. The refined deodorized sardine oil (distributed by Mochida International Company, Limited, Shinjuku-ku, Tokyo, Japan under the trade designation "50% Omega-3 marine oil EPA:DHA 28:12 with 0.8% mixed tocopherol as antioxidant") is then added to the slurry with agitation. (In a most preferred method of manufacture the sardine oil would be slowly metered into the product as the blend passes through a conduit at a constant rate.) Preferably after at least 5 minutes the pH of the blended slurry is determined. If the pH of the blended slurry is below 6.55, it is adjusted with dilute potassium hydroxide to a pH of 6.55 to 6.8.

After waiting a period of not less than one minute nor greater than two 45 hours the blended slurry is subjected to deaeration, Ultra-High-Temperature (UHT) treatment, and homogenization, as described below:

A. Use a positive pump for supplying the blended slurry for this procedure.

- B. Heat the blended slurry to a temperature in the range of about 66 - 71 °C.
- C. Deaerate the blended slurry to 25.4 - 38.1 cm of Hg.
- D. Emulsify the blended slurry at 61-75 Atmospheres.
- E. Heat the blended slurry to a temperature in the range of about 120 - 122 °C
- 5 by passing it through a plate/coil heat exchanger with a hold time of approximately 10 seconds.
- F. UHT heat the blended slurry to a temperature in the range of about 144 - 147 °C with a hold time of approximately 5 seconds.
- G. Reduce the temperature of the blended slurry to be in the range of about
- 10 120-122 °C by passing it through a flash cooler.
- H. Reduce the temperature of the blended slurry to be in the range of about 71 - 82 °C by passing it through a plate/coil heat exchanger.
- I. Homogenize the blended slurry at about 265 to 266 Atmospheres.
- J. Pass the blended slurry through a hold tube for at least 16 seconds at a
- 15 temperature in the range of about 74 - 85 °C.
- K. Cool the blended slurry to a temperature in the range of about 1 - 70 °C by passing it through a large heat exchanger.

Store the blended slurry at a temperature in the range of about 1 - 7 °C, preferably

20 with agitation.

Preferably at this time appropriate analytical testing for quality control is conducted. Based on the test results an appropriate amount of dilution water (10-38 °C) is added to the blended slurry with agitation.

A vitamin solution, a flavor and a color solution are prepared separately and then

25 added to the blended slurry.

The vitamin solution is prepared by heating about 394 kg of water to a temperature in the range of about 43 - 66 °C with agitation, and thereafter adding the following ingredients, in the order listed: Ascorbic Acid, 45% Potassium Hydroxide, Taurine, Water Soluble Vitamin Premix, Folic Acid, Choline Chloride, and L-Carnitine. The vitamin

30 solution is then added to the blended slurry with agitation.

The flavor solution is prepared by adding the artificial strawberry flavor and artificial cream flavor to about 794 kg of water with agitation. A nutritional product according to the present invention has been manufactured using an artificial strawberry flavor distributed by Firmenich Inc., Princeton, New Jersey, U.S.A. under the trade

designation "Art. strawberry 57.883/A" and an artificial cream flavor distributed by Firmenich Inc. under the trade designation "Art Cream 59.200/A". The flavor solution is then added to the blended slurry with agitation.

A color solution is prepared by adding the FD&C Red Dye No. 3 to about 121 kg of water with agitation. The color solution is then added to the blended slurry with agitation.

If necessary, diluted potassium hydroxide is added to the blended slurry such that the product will have a pH in the range of 6.4 to 7.0 after sterilization. The completed product is then placed in suitable containers and subjected to sterilization. Of course, if desired aseptic processing could be employed.

The product made according to the procedure of this example contains the oil blend of Table 8, below, the fatty acid properties of Tables 2 and 3, and the amino acid profile of Table 4, all set forth above.

TABLE 8: OIL BLEND

(% total weight of lipid blend)

<u>OIL</u>	<u>Percent Total Lipids</u>
Canola	9.3%
MCT	16.2%
Fish	65.0%
Soybean	5.5%
Soy lecithin	4.0%

EXAMPLE II

The objective of this experiment was to evaluate the organoleptic characteristics of nutritional composition of the invention fortified by the addition of branched-chain amino acids incorporated at two different levels. To measure organoleptic properties, three taste standards, described in Table 9, were prepared to rank the bitter and sour intensity of the test compositions containing branched-chain amino acids.

TABLE 9: TASTE INTENSITY SCALE

Standard	Basic Taste	Intensity	Concentration * by Weight	Representative Products
1	Sour	1	0.05% Citric Acid	Milk Chocolate, Coffee
2	Bitter	1	0.05% Caffeine	Whole Peanuts Milk Chocolate
3	Bitter	2	0.10% Caffeine	Beer

* Aqueous solutions

Two test compositions (designated "high" and "low") were prepared by adding selected branched-chain amino acids ("BCAA") to the liquid nutritional composition of Example I. A control composition of Example I that did not contain the supplemental branched-chain amino acids was also evaluated for flavor characteristics. The specific amino acids and amounts added are given in Table 10. In both test compositions the branched-chain amino acids did not completely disperse in the nutritional composition due to their hydrophobic nature, and small clumps of branched-chain amino acids were visible in the matrix. The test compositions were evaluated and the results of the organoleptic test scoring are also set forth in Table 10.

TABLE 10: BRANCHED-CHAIN AMINO ACID FORTIFICATION

	"high" test	"low" test	control
BCAA in gm/237 mL serving			
valine	2.5	1.3	0
leucine	2.5	1.3	0
<u>isoleucine</u>	<u>2.5</u>	<u>1.3</u>	<u>0</u>
Total	7.5	3.9	0
TASTE TEST SCORE			
bitter	1.5 to 2	1.5	none
sour	0.5	0.5	none

15

Based on the results of this taste session, the evaluators collectively agreed that the bitter and sour flavor notes attributed to the branched-chain amino acids are less than ideal for a ready-to-use oral nutritional composition. Thus, in one embodiment of this invention, any additional branched-chain amino acids are supplied to patient in the form of a pill or capsule distinct from the liquid nutritional of the invention..

20

EXAMPLE III

The effect of nutritional intervention with ω -3 fatty acids, branched-chain amino acids and antioxidants in the nutritional compositions of the invention, on prevention and treatment of cachexia can be monitored by any of the methods known to one skilled in the art, including but not limited to measuring: (i) food intake, body weight and anthropometric measurements; (ii) serum levels of lipids, fatty acids, amino acids and antioxidants; (iii) levels of serologic markers where appropriate, e.g., carcinoembryonic (CEA) antigens, serotonin, C-reactive protein, TNF and IL-1; (iv) changes in the morphology of tumors

25

using techniques such as computed tomographic (CT) scan, ultrasonography, magnetic resonance imaging (MRI) and position emission tomography (PET).

Patients with hepatocellular carcinoma showing symptoms of cachexia are provided with the nutritional product of the invention with small, frequent feedings after surgical resection if the liver tumor is localized and small, or along with a regimen of chemotherapy. The liver functions and characteristics of the hepatic carcinomas are tested by procedures known in the art.

The daily nutritional management of liver cancer, therefore, includes administration of 2 to 4 containers of 8 ounce servings (237 mL) of the nutritional composition providing a daily amount of: (i) combined EPA and DHA in the range of 3 to 6g, with the preferred dosage being about 3g; (ii) branched-chain amino acids in the range of 5 to 25 g, with the preferred dosage being about 10-15g branched-chain amino acids; and (iii) vitamin C in the range of 125 to 500 mg, with the preferred dosage being about 300 mg vitamin C; (iv) vitamin E in the range of 50 to 250 IU, with the preferred dosage being about 150 IU vitamin E; (v) beta-carotene in the range of 1250 to 3250 µg, with the preferred dosage being about 2500 beta-carotene µg; (vi) selenium in the range of 40 to 60 µg, with the preferred dosage being about 45 µg selenium. The effect of nutritional intervention on cancer cachexia and anorexia are monitored at monthly intervals (or as recommended in the clinical follow-up) as known in the art, and depending on the results obtained, the therapeutic regimen is developed to maintain and/or boost the weight gain by the patient, with the ultimate goal of achieving tumor regression and complete eradication of cancer cells.

For the underweight breast cancer patient on adjuvant chemotherapy, administration of the nutritional composition of the invention is started any time after surgery. The nutritional composition used in breast cancer patients is designed to maintain an adequate intake in spite of nausea, mucositis, and other side effects of chemotherapy. Patients receiving radiation therapy for breast cancer receive effective amounts of the nutritional composition to promote maintenance and repair of body tissue. The therapeutic and/or prophylactic regimens used in breast cancer patients are the same as those described in Section 6 above for patients recovering from hepatocellular carcinoma. The procedures of monitoring the patient under clinical evaluation for prevention and treatment of cachexia and anorexia in breast cancer are known in the art.

EXAMPLE IV

A pilot study was conducted to assess the effectiveness of a specific formula in ameliorating the cachexia of cancer patients. The formula of Example I was prepared. In addition to other nutrients, it contained (per two 237 mL servings) the long-chain ω -3 fatty acids, the fructooligosaccharide ("FOS") and the antioxidant system specified in Table 10.

TABLE 10: TRIAL PRODUCT

10	Ingredient	Amount per 2 x 237 mL servings
	EPA ω -3	2.0 gm
	DHA ω -3	0.92 gm
	fructooligosaccharide	5.8 gm
15	beta carotene	2.8 mg
	vitamin C	300. mg
	vitamin E	150. IU
	selenium	58. mcg

In the pilot clinical trial of this example, ten patients with pancreatic cancer were evaluated. These patients were cachectic and losing weight at a mean rate of 0.86 kg per week over an average of 22 weeks (range: 11 to 56 weeks) prior to the trial. Over a three week trial period, patients consumed an average of two 237 mL (8 fluid ounces) servings per day as a supplement to their diets. After the trial period the group demonstrated a mean increase in weight of 2.1 kg (up from baseline), which translates to a mean weekly weight gain of 0.7 kg (See Table 11).

TABLE 11: Patients' Age, Gender and Weight Status

Patient	Patient Age and Gender	Mean Weekly Wt (kg) Change up to Baseline	Baseline weight (kg)	Weight after three week trial (kg)	Mean Weekly Wt change during trial
1	56f	-0.4	51	51.75	0.25
2	64m	-1.2	67	na	na
3	70m	-0.5	61	67.5	2.17
4	60m	-0.9	43	44	0.33
5	53f	-1.4	90	90.5	0.17
6	51f	-1.2	44.5	45.5	0.33
7	67m	-0.6	57	58	0.33
8	75f	-0.8	55	59	1.33
9	57m	-0.6	69	na	na
10	53f	-1	57.5	na	na
Mean		-0.86			0.7

In addition, much of the weight gained was lean body mass. The group demonstrated a mean increase in lean body mass of 2.1% and a decrease in C-reactive protein ("CRP") levels (See Table 12). Serum CRP is a biochemical surrogate for the presence and progress of cancer cachexia, and shows a strong positive correlation. (Falconer, J.S. et al., *Cancer* 1995, 75:2077). Patients with serum CRP levels >10 mg/L are frankly cachectic. The mean CRP level at baseline was 31.5 and this dropped to about 10 after 3 weeks on the experimental formula of the invention. Thus, the invention improves cachexia in pancreatic cancer patients.

10 **TABLE 12: Patients' Lean Body Mass and CRP**

Patient	% Lean Body Mass at Baseline	% Lean Body Mass after trial	Change in % Lean Body Mass	Baseline CRP mg/L	CRP after 3 week trial mg/L
1	79.4	82.9	3.5	69	<10
2	85.6	na	na	81	na
3	84	92.2	8.2	27	<10
4	85.6	86.3	0.7	<10	<10
5	73.7	75	1.3	<10	<10
6	79.7	80.7	1	63	<10
7	90.2	91.9	1.7	<10	10
8	81.4	80.1	-1.3	<10	<10
9	82.6	na	na	25	na
10	86.9	na	na	<10	na
Mean	82.9	84.2	2.1	31.5	10*

* CRP values read as <10 are assumed to be 10 for calculation of mean

The present invention is not to be limited to the scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

Various publications and patents are cited herein, the disclosures of which are incorporated by reference in their entireties.

We claim:

1. A method of preventing the onset of cachexia and/or anorexia or treating existing cachexia and/or anorexia in a human, said method comprising enterally administering to the human on a daily basis at least:

(a) an oil blend containing ω -6 fatty acids and at least 450 mg of ω -3 fatty acids, the weight ratio of ω -6 fatty acids to ω -3 fatty acids being from about 0.1 to about 3.0; and

(b) a source of amino-nitrogen wherein 15% to 50% by weight of the amino acids of said source of amino-nitrogen are branched-chain amino acids; and

(c) an antioxidant component comprising at least one nutrient selected from the group comprising beta-carotene, vitamin C, vitamin E, selenium, and mixtures thereof;

wherein at least said oil blend and said antioxidant are present in a liquid nutritional matrix.

2. The method according to claim 1, further comprising administering to the human a source of amino-nitrogen wherein tryptophan is present in an amount of less than about 5.0% by weight of the total amount of the amino acids present in said source of amino-nitrogen.

3. The method according to claim 1, further comprising administering to the human a source of amino-nitrogen wherein 15% to 25% by weight of the amino acids of said source of amino-nitrogen are branched-chain amino acids.

4. The method according to claim 1 which further comprises daily administering to the human:

(a) an oil blend containing ω -6 fatty acids and at least 1000 mg of ω -3 fatty acids, the weight ratio of ω -6 fatty acids to ω -3 fatty acids being from 0.1 to 1.0; and

(b) a source of amino-nitrogen wherein 15% to 25% by weight of the amino acids are branched-chain amino acids, and wherein tryptophan is present in an amount of less than 3.0% by weight of the total amount of the amino acids present in said source of amino-nitrogen; and

(c) an antioxidant mixture comprising beta-carotene, vitamin C, vitamin E and selenium.

5. The method of claim 4 wherein the antioxidant system of component (c) comprises about 2,500 to about 6,500 micrograms beta-carotene, about 250 to about 1,000 milligrams vitamin C, about 100 to about 500 I.U. vitamin E and about 75 to about 125 micrograms selenium per liter.
6. The method of claim 1 in which the ω -3 fatty acids comprise from about 15 to about 30% by weight of the total fatty acids and are selected from the group consisting of alpha-linolenic acid, stearidonic acid, eicosapentaenoic acid, docosapentaenoic acid and docosahexaenoic acid.
7. The method of claim 6 in which the ω -3 fatty acids comprise from about 15 to about 40% by weight of the total fatty acids and are selected from the group consisting of eicosapentaenoic acid and docosahexaenoic acid.
8. The method of claim 7 in which at least about 15% by weight of the total fatty acids is eicosapentaenoic acid.
9. The method of claim 1 wherein the weight ratio of ω -6 fatty acids to ω -3 fatty acids in said oil blend is from about 0.1 to about 1.0; about 15% to about 25% by weight of the amino acids of said source of amino-nitrogen are branched-chain amino acids; and said antioxidant component comprises a mixture of beta-carotene, vitamin C, vitamin E and selenium.
10. The method according to claim 1 which further comprises daily administering to the human:
 - (a) an oil blend containing ω -6 fatty acids and at least 1000 mg of ω -3 fatty acids, the weight ratio of ω -6 fatty acids to ω -3 fatty acids being from 0.1 to 1.0; and
 - (b) a source of amino-nitrogen wherein 15% to 25% by weight of the amino acids are branched-chain amino acids, and wherein tryptophan is present in an amount of less than 5.0% by weight of the total amount of the amino acids present in said source of amino-nitrogen.
11. The method of claim 10 comprising feeding eicosapentaneic acid as the ω -3 fatty acid.

12. A liquid nutritional composition comprising per liter:
 - (a) at least 450 mg of ω -3 fatty acids and wherein the weight ratio of ω -6 fatty acids to ω -3 fatty acids is from about 0.1 to about 3.0;
 - (b) at least 50 grams of a source of amino-nitrogen wherein 15 to 50% by weight of the amino-nitrogen is branched-chain amino acids and wherein tryptophan is present in an amount less than about 5.0% by weight of the total amino-nitrogen; and
 - (c) at least 1 gram of an antioxidant system comprising beta-carotene, vitamin C, vitamin E and selenium.
13. The composition according to claim 12, wherein the weight ratio of ω -6 fatty acids to ω -3 fatty acids is from about 0.1 to about 1.0, and wherein 15 to 25% by weight of the amino-nitrogen is branched-chain amino acids.
14. The composition according to claim 13, further comprising at least 1000 mg of ω -3 fatty acids per liter.
15. A liquid nutritional composition according to claim 12 wherein:
 - (a) at least 500 mg of ω -3 fatty acids are present and the weight ratio of ω -6 fatty acids to ω -3 fatty acids is from 0.2 to 2.0;
 - (b) tryptophan is present in an amount of less than 3.0% by weight of the total amino-nitrogen; and
 - (c) at least 2.0 grams of said antioxidant system is present per liter.
16. A method of preventing the onset of anorexia or of treating existing anorexia in a human comprising administering to the human a nutritional composition comprising amino acids wherein about 5 to 25 grams of branched-chain amino acids selected from valine, leucine, isoleucine, or mixtures thereof are present in an amount from about 15% to about 50% by weight of the total amount of amino acids present in said composition, and wherein tryptophan in an amount not greater than about 5.0 % by weight of the total amount of amino acids is present in said composition and wherein ω -6 and ω -3 fatty acids are present at a weight ratio of from about 0.1 to about 3.0 and at least one antioxidant is present in the nutritional composition.

17. A method for preventing immunosuppression in a human comprising administering to the human a liquid nutritional composition comprising:

- (a) an oil blend containing ω -6 and ω -3 fatty acids, the weight ratio ω -6 fatty acids to ω -3 fatty acids being about from 0.1 to about 3.0; and
- (b) an antioxidant component comprising about 2,500 to about 6,500 micrograms per liter beta-carotene, about 250 to about 1,000 milligrams per liter vitamin C, about 100 to about 500 I.U. per liter vitamin E, and about 75 to about 125 microgram per liter selenium.

18. The method of claim 17 in which at least 400 mL of said liquid nutritional composition are administered to said human per day.

19. A method for enhancing the transport and efficacy of anticancer drugs in a human having a cancerous condition comprising administering to the human a nutritional composition comprising an oil blend containing ω -6 and ω -3 fatty acids, the weight ratio of total ω -6 fatty acids to ω -3 fatty acids being from about 0.1 to about 3.0.

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/20 // (A61K 31/20, 31:195, 31:00)		A3	(11) International Publication Number: WO 97/39749
			(43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US97/06897		(81) Designated States: AU, CA, JP, MX, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 25 April 1997 (25.04.97)		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 08/635,179 25 April 1996 (25.04.96) US 08/842,454 24 April 1997 (24.04.97) US		(88) Date of publication of the international search report: 4 December 1997 (04.12.97)	
(71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).			
(72) Inventors: ABBRUZZESE, Bonnie, Chandler; 5673 Keating Drive, Dublin, OH 43017 (US). McCAMISH, Mark, Anthony; 1084 Woodman Drive, Worthington, OH 43085 (US). COPE, Frederick, Oliver; 6826 McGregor Street, Worthington, OH 43085 (US). DEMICHELE, Stephen, Joseph; 5525 Windwood Drive, Dublin, OH 43017 (US).			
(74) Agents: BRAINARD, Thomas, D.; Abbott Laboratories, Ross Products Division, Dept. 108140/DS-1, 625 Cleveland Avenue, Columbus, OH 43215-1724 (US) et al.			
(54) Title: METHOD FOR THE PREVENTION AND TREATMENT OF CACHEXIA AND ANOREXIA			
(57) Abstract The present invention relates to methods and nutritional compositions for the prevention and treatment of cachexia and anorexia. The methods of the invention comprise administering a composition comprising effective amounts of ω -3 fatty acids such as alpha-linolenic acid, stearidonic acid, eicosapentaenoic acid, docosapentaenoic acid, docosahexaenoic acid or mixtures thereof; of branched-chain amino acids valine, leucine, isoleucine or mixtures thereof; with or without reduced levels of tryptophan and 5-hydroxytryptophan; and of antioxidant system selected from the group comprising beta-carotene, vitamin C, vitamin E, selenium, or mixtures thereof.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/06897

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/20 //(A61K31/20,31:195,31:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 97 13415 A (COLGATE PALMOLIVE CO ;OGILVIE GREGORY K (US)) 17 April 1997 see page 8-12 see claims 1-9	1-15
Y	--- ANTICANCER RESEARCH, vol. 14, 1994, pages 1451-1455, XP002035801 C. CANGIANO ET AL: "CYTOKINES, TRYPTOPHAN AND ANOREXIA IN CANCER PATIENTS BEFORE AND AFTER SURGICAL TUMOR ABLATION" cited in the application see the whole document	1-15
Y	--- EP 0 642 791 A (NEW YORK STATE OFFICE OF MENTA) 15 March 1995 see the whole document --- -/--	1-15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

23 September 1997

Date of mailing of the international search report

24. 10. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Herrera, S.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/06897

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 367 724 A (SANDOZ AG ;SANDOZ AG (DE); SANDOZ AG (AT)) 9 May 1990 see abstract; claims 1-4 ---	17,18
P,X	EP 0 756 827 A (BRAUN MELSUNGEN AG) 5 February 1997 see abstract see page 3, column 58 - page 4, column 14 ---	17,18
Y	US 5 308 832 A (GARLEB KEITH A ET AL) 3 May 1994 see abstract ---	17,18
X	WO 95 08349 A (FLOCKHART IAN ;ALTUNKAYA ALI (GB); HOPEWELL JOHN WILLIAM (GB)) 30 March 1995 see abstract ---	19
X	EP 0 462 398 A (HOFFMANN LA ROCHE) 27 December 1991 see abstract -----	19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/06897

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Claims 1-15
Claim 16
Claims 17-18
Claim 19

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

17-18 and 19
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inten. nal Application No

PCT/US 97/06897

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9713415 A	17-04-97	AU 7254596 A	30-04-97
EP 0642791 A	15-03-95	US 5393784 A	28-02-95
		CA 2128486 A	22-01-95
EP 0367724 A	09-05-90	AU 626930 B	13-08-92
		AU 4386089 A	03-05-90
		CA 2001727 A	30-04-90
		ES 2052066 T	01-07-94
		HK 34694 A	22-04-94
		JP 2191213 A	27-07-90
		JP 2520488 B	31-07-96
		SG 30494 A	10-06-94
		US 5231085 A	27-07-93
EP 0756827 A	05-02-97	DE 19528461 A	06-02-97
		JP 9121809 A	13-05-97
US 5308832 A	03-05-94	AU 666246 B	01-02-96
		AU 5574794 A	14-02-94
		EP 0656782 A	14-06-95
		JP 7507327 T	10-08-95
		MX 9304468 A	28-02-94
		WO 9402166 A	03-02-94
WO 9508349 A	30-03-95	AU 7787194 A	10-04-95
		EP 0755267 A	29-01-97
EP 0462398 A	27-12-91	US 5147854 A	15-09-92
		AU 636489 B	29-04-93
		AU 7713991 A	12-12-91
		CA 2042973 A	23-11-91
		JP 5213772 A	24-08-93

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)